

NOTE

**DRY MATTER, ENERGY, AND NUTRIENT CONTENTS
OF 8-YEAR-OLD STANDS OF
EUCALYPTUS REGNANS, *ACACIA DEALBATA*,
AND *PINUS RADIATA* IN NEW ZEALAND**

D. J. FREDERICK, H. A. I. MADGWICK, M. F. JURGENSEN, AND G. R. OLIVER

Ministry of Forestry, Forest Research Institute, Private Bag 3020,
Rotorua, New Zealand

(Received for publication 18 December 1990)

An error in energy content units has recently been discovered in Table 4 of the original paper (Frederick *et al.* 1985). The corrected version is presented below.

TABLE 4—Average calorific value and energy content of 8-year-old stands of *A. dealbata*, *E. regnans*, and *P. radiata*

	<i>A. dealbata</i>		<i>E. regnans</i>		<i>P. radiata</i>	
	kJ/g	10 ¹¹ J/ha	kJ/g	10 ¹¹ J/ha	kJ/g	10 ¹¹ J/ha
Leaves	21.8	1.4	24.4	2.1	21.2	1.9
Live branches	19.3	2.8	19.4	3.1	20.1	4.6
Dead branches	19.8	3.1	19.4	2.4	21.5	0.5
Fruits	—	0.0	—	0.0	20.0	0.1
Stem wood	19.3	25.0	19.4	23.6	20.4	12.5
Stem bark	19.9	3.1	17.6	2.2	20.4	1.5
Tree above-ground	—	35.4	—	33.5	—	21.1

REFERENCE

- FREDERICK, D.J.; MADGWICK, H.A.I.; JURGENSEN, M.F.; OLIVER, G.R. 1985: Dry matter, energy, and nutrient contents of 8-year-old stands of *Eucalyptus regnans*, *Acacia dealbata*, and *Pinus radiata* in New Zealand. *New Zealand Journal of Forestry Science* 15: 142–57.